

VULVAR LICHEN PLANUS



What is it?

Lichen planus (LP) is a disease of the skin caused by inflammation. Vulvar lichen planus occurs most commonly in women 50-60 years old. It can affect the genital area including both the vulva and the vagina. The most common symptoms are burning and soreness. Some women describe itching as well. Lichen planus causes a rash of small purplish bumps, often on the arms, legs, or back. It can affect the mouth (oral disease) with a whitish pattern or loss of the mouth surface. In some cases, the nails and the scalp are also involved. It is possible to have the disease in one area without ever having a problem elsewhere. Many patients with vulvovaginal LP have LP in the mouth as well and sometimes on other areas of the skin.

How common is it?

It is estimated that 1 in 4000 women will have vulvar or vaginal LP compared to 1 in 100 who may have oral LP. About 50% of women who have oral lichen planus may have vulvar or vaginal LP but the diagnosis may be missed as dentists do not generally enquire about genital symptoms and the mouth may not be routinely examined in those presenting with genital disease.

What causes LP?

The cause of LP is unknown. There may be a problem with the immune system, the system that protects a person from diseases. In LP, the system is overactive and can act against itself (this is called an auto-immune reaction). In some cases, it is possible that an infection or medication can start this reaction. We do not know why the lesions develop in some parts of the body and not others. LP may be associated with other auto-immune conditions such as thyroid disease, vitiligo (white patches on the skin), and alopecia areata (patches of hair loss) Lichen planus is NOT infectious or contagious and cannot be passed to a sexual partner or to another part of your body.

What are the symptoms and what do I see?

Soreness, burning and rawness are very common symptoms. Less commonly, itching is present. If the outer layers of the skin break down (erosions), these areas appear moist and red. There may be a white lacy pattern on the vulva. This pattern can also be seen around the edges of the erosions. The vulva may appear pale white or pink/red. Scarring with loss of the inner lips (labia minora) may be seen. The clitoris may be buried under scar tissue. There may be shiny, red, raw areas. Intercourse can be painful if the vagina is involved or there is scar tissue narrowing the entrance of the vagina. Erosions can occur inside the vagina in a patchy or generalized pattern. Some women have a sticky, yellow or yellow-green discharge, which can be bloodstained, especially after intercourse. The vaginal entrance may become smaller if the inner walls of the vagina or the skin around the entrance stick together when it heals.

This is one reason why intercourse can be painful or even impossible. Sometimes it is difficult for a healthcare provider to perform an internal examination. On rare occasions, the skin may have thickened areas. These may have a warty appearance. If the skin is affected in other parts of the body, the rash is usually on the inside of the wrist, the forearms and the ankles. The spots are a purple color and you may see some fine white streaks on the top of the spots. A similar white, lacy streaking may be seen inside the mouth, but there may not be any symptoms. There may be sore, red, ulcerated areas around the gum margins, tongue and inside of cheeks. Occasionally LP can affect the tear ducts and esophagus (the tube that carries food from the mouth to the stomach). If you experience excessive watering of the eyes, difficulty in swallowing or it feels as if food gets stuck, you should tell your doctor about this.

What should I watch for?

As LP is a long lasting inflammatory skin condition, there is an increased risk of developing local types of skin cancer in the area compared to women without LP. The risk is about 3 percent. Any new raised, bleeding or non-healing areas in your genital area should be reported to your healthcare provider.

It is important that your LP is monitored and that you attend for follow-up visits with your healthcare provider at least once per year.